



This is a repository copy of *The exosome-binding factors Rrp6 and Rrp47 form a composite surface for recruiting the Mtr4 helicase.*

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/89310/>

Version: Supplemental Material

---

**Article:**

Schuch, B., Feigenbutz, M., Makino, D.L. et al. (4 more authors) (2014) The exosome-binding factors Rrp6 and Rrp47 form a composite surface for recruiting the Mtr4 helicase. EMBO Journal, 33 (23). 2829 - 2846. ISSN 0261-4189

10.15252/emj.201488757

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

